arXiv:hep-ph/0211047v2 6 Nov 2002

Measurement of the Higgs self-coupling at JLC

Y. Yasui[1†*], S. Kanemura[1†], S. Kiyoura[1,2†], K. Odagiri[1†], Y. Okada[1‡], E. Senaha[1,3]
and S. Yamashita[4§]

[1] *KEK, Tsukuba,1-1 Oho, Tsukuba, Ibaraki 305-0801,Japan*
[2] *Department of Radiological Sciences, Ibaraki Prefectural University of Health Sciences, Ami, Inashiki, Ibaraki 300-0394, Japan*
[3] *Department of Particle and Nuclear Physics, the Graduate University for Advanced Studies, Tsukuba, Ibaraki 305-0801,Japan*
[4] *ICEP, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, JAPAN*

Abstract

We examine the double Higgs production process at JLC. We focus our attention on the measurement of the Higgs self-coupling. The sensitivity of the triple Higgs coupling measurement is discussed in the Higgs mass range 100-200 GeV and the center of mass energy to be 500 GeV-1.5 TeV.

After the discovery of the Higgs boson, detailed study of the Higgs potential will be one of the important issues at the linear collider (LC) experiment[1, 2]. From the standard model (SM) Higgs potential $V(\Phi) = -\mu^2\Phi^\dagger\Phi + \lambda(\Phi^\dagger\Phi)^2$, we obtain the simple relation between the Higgs boson mass m_h and the self coupling λ as $m_h^2 = 2\lambda v^2$, where v is the vacuum expectation value of the Higgs boson. The precision test of this relation could give clear evidence for the new physics[3]. At the LC, multiple Higgs boson production processes will provide the direct information of the triple Higgs coupling. In this talk, we discuss the double Higgs production at JLC[4]. We examine the potential of the triple Higgs coupling measurement.

At JLC, we can use the two modes, $e^+e^- \to hhZ$ and $e^+e^- \to (WW)\nu\bar{\nu} \to hh\nu\bar{\nu}$. In the low center of mass energy region($\sqrt{s} \lesssim 1$ TeV), the hhZ mode is dominant. On the other hand, the $hh\nu\bar{\nu}$ mode will be important at the 1 TeV or higher energies. Here, we introduce the effective coupling which includes the anomalous contribution

---

[*] e-mail address:yoshiaki.yasui@kek.jp; This work was supported in part by Grand-in-Aid for Scientific Research [†](A)(1) (No. 13047101), [‡](C) (No. 13640309), [§](A)(No.14046205).

as $V(h^3) = (\lambda_3 + \delta\lambda_3)hhh$, where $\lambda_3(\equiv \lambda v)$ is the SM triple Higgs coupling and $\delta\lambda_3$ is the deviation from the SM prediction. In figure 1, we show the $\delta\lambda_3$ dependence of the total cross section for (a) $e^+e^- \to hhZ$ and (b) $e^+e^- \to hh\nu\bar{\nu}$. Here, the Higgs mass is 120 GeV with the CM energy of 500 GeV, 1 TeV and 1.5 TeV. To compute the cross section, we mainly employ the GRACE[5] system, and use the CompHEP[6] system for the cross-check. In the low values of the invariant mass M_{hh} for hh, the hhZ mode is sensitive to $\delta\lambda_3$(Fig. 2). We found that the cut for M_{hh} is powerful to improve the sensitivity. The polarized electron beam is also useful to reduce the back-ground for the $hh\nu\bar{\nu}$ mode, due to the $V - A$ nature of the weak bosons.

We show the sensitivity of the triple Higgs coupling to $\delta\lambda_3$ in the mass range $100 - 200$ GeV at the parton level (Fig. 3). Here, we assumed that the efficiency of the particle tagging is 100% with an integrated luminosity of 1 ab^{-1}. Dashed (dotted, sold) lines are the results of the hhZ mode ($hh\nu\bar{\nu}$ mode, combined both modes). At $\sqrt{s} = 500$ GeV, the hhZ mode is dominant. If the Higgs mass is relatively light ($m_h \lesssim 150$ GeV), we can measure the triple Higgs coupling in about 20% accuracy. For $\sqrt{s}$ to be 1 TeV or higher, the $hh\nu\bar{\nu}$ mode is dominant. Here, we used the invariant mass cut as $M_{hh} < 600$ GeV for hhZ mode. We also used the 100% polarized electron beam for the $hh\nu\bar{\nu}$ mode. We can expect the high sensitivity in this case ($\delta\lambda_3/\lambda_3 \lesssim 10\%$).

Acknowledgment We would like to thank Y. Kurihara, T. Kaneko, T. Ishikawa, J. Fujimoto, Y. Shimizu and S. Kim for valuable discussions.


(a) (b)

Figure 1: The $\delta\lambda_3$ dependence of the cross section; (a) $e^+e^- \to hhZ$, (b)$e^+e^- \to hh\nu\bar{\nu}$



Figure 2: The hh imvariant mass dependence of the hhZ mode for several values of $\delta\lambda_3$.



Figure 3: The λ_3 measurement sensitivity; hhZ (dashed line), $hh\nu\bar{\nu}$ (dotted line) and combined results (solid line).

References

[1] F.Boudjema and E. Chopin, **Z.Phys. C73** (1996) 85; V.A.Ilyin, T.Kaneko, Y.Kurihara, A.E.Pukhov and Y.Shimizu, **Phys.Rev. D54** (1996) 6717; Marco Battaglia and Klaus Desch, "Studying the Higgs Potential at the e+e- Linear Collider" Proceedings of the Linear Collider Workshop LCWS2000; arXiv:hep-ph/0111276.

[2] A. Djouadi, W. Kilian, M. Muhlleitnera and P.M. Zerwas, **Eur.Phys.J. C10** (1999) 45.

[3] See for example, the SUSY model was discussed in [2], the 2HDM was discussed in S. Kanemura, S. Kiyoura, Y. Okada, E. Senaha and C.-P. Yuan, Proceedings of the Linear Collider Workshop LCWS2002; arXiv:hep-ph/0209326.

[4] ACFA Linear Collider Working Group report, KEK Report 2001-11; arXiv:hep-ph/0109166.

[5] J. Fujimoto, T. Ishikawa, M. Jimbo, T. Kaneko, K. Kato, S. Kawabata, K. Kon, M. Kuroda, Y.Kurihara, Y. Shimizu and H. Tanaka, KEK-CP-129; arXiv:hep-ph/0208036.

[6] A.Pukhov, E.Boos, M.Dubinin, V.Edneral, V.Ilyin, D.Kovalenko, A.Kryukov, V.Savrin, S.Shichanin, and A.Semenov, INP MSU 98-41/542; arXiv:hep-ph/9908288